Centre Number	Candidate Number	Name

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CHEMISTRY 0620/06

Paper 6 Alternative to Practical

May/June 2005

1 hour

Candidates answer on the Question Paper. No additional materials required.

READ THESE INSTRUCTIONS FIRST

Write your name, Centre number and candidate number at the top of this page. Write in dark blue or black pen in the spaces provided on the Question Paper. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid.

Answer all questions.

The number of marks is given in brackets [] at the end of each question or part question.


If you have been given a label, look at the details. If any details are incorrect or missing, please fill in your correct details in the space given at the top of this page.

Stick your personal label here, if provided.

FOR EXAM	INER'S USE
1	
2	
3	
4	
5	
6	
7	
8	
TOTAL	


[1]

1 A small piece of limestone was heated strongly and left to cool. A few drops of cold water were added. The solid expanded and gave off steam.




- (a) Complete the empty boxes to identify the pieces of apparatus labelled. [3]
- (b) What type of chemical reaction takes place when water is added?

2 The diagram shows the apparatus used to find out the effect of an electric current on a concentrated aqueous solution of sodium chloride.



(a)	On	the diag	gram label the electrodes	[1]
(b)	Giv	e three	observations when the circuit is switched on.	
	1			
	2			
	3			[3]
(c)	(i)	Name	the product at the positive electrode (anode).	[1]
	(ii)	State a	a test for this product and the result of the test.	ניו
		test		
		result		[2]

3 In a set of experiments zinc was reacted with sulphuric acid to form hydrogen. The apparatus below was used.




The same mass of zinc was used each time. The volume of acid used was different each time. Use the syringe diagrams to record the volume of hydrogen produced each time in the table.

Table of results

volume of sulphuric acid/cm ³	syringe diagram	volume of hydrogen/cm ³
0	10 20 30 40 50 60 70 80	
5	10 20 30 40 50 60 70 80	
15	10 20 30 40 50 60 70 80	
20	10 20 30 40 50 60 70 80	
25	10 20 30 40 50 60 70 80	
30	10 20 30 40 50 60 70 80	
35	10 20 30 40 50 60 70 80	
40	10 20 30 40 50 60 70 80	

(a) Plot the results on the grid below. Draw a smooth line graph.




(c) What volume of gas is produced if 10 cm³ of sulphuric acid is used? [1] 4 A student investigated an aqueous solution of calcium hydroxide and water.

Two experiments were carried out.

Experiment 1

By using a measuring cylinder 25 cm³ of the aqueous solution of calcium hydroxide was placed in a flask. Phenolphthalein indicator was added to the flask. A burette was filled to the 0.0 cm³ mark with solution **M** of hydrochloric acid.


Solution **M** was added slowly to the flask until the colour just disappeared. Use the burette diagram to record the volume in the table and complete the column.




Experiment 2

Experiment 1 was repeated using a different solution, **N**, of hydrochloric acid.

Use the burette diagrams to record the volumes in the table and complete the table.





final

Table of results

burette readings/cm ³	Experiment 1	Experiment 2
final reading		
initial reading	0.0	
difference		

			4]
(a)		at type of chemical reaction occurs when hydrochloric acid reacts with calciudroxide?	ım
		[[1]
(b)	(i)	In which experiment was the greater volume of hydrochloric acid used?	
			[1]
	(ii)	Compare the volumes of acid used in Experiments 1 and 2.	
		ı	[2]
			, - _1
	(iii)	Suggest an explanation for the difference in volumes.	
		ı	[2]
			,
(c)		edict the volume of hydrochloric acid M that would be needed to react completely periment 1 was repeated with 50 cm ³ of calcium hydroxide solution?	' if
	vol	ume of solution	
	exp	olanation	
			[3]
(d)		ggest one change you could make to the apparatus used in the experiments ain more accurate results.	to
		[[1]

5 A sample of a solution of acid **A** was analysed.

The tests on **A**, and some of the observations are in the following table.


Complete the observations in the table.

		tests			observations	
(a)		e pH of the solution w sted using indicator p		colour	orange	
	ıc	sted using indicator p		рН	4	
(b)		e solution was divided ee test-tubes	l into			
	(i)	To the first portion wadded a piece of magnesium ribbon. gas was tested with lighted splint.	Гће			[2
	(ii)	To the second portion A was added sodium carbonate. The gas tested with limewate	n was			[2
	(iii)	To the third portion of liquid A was added a spatula measure of B . The mixture was gently. By using a tepipette the solution of transferred to another.	a solid boiled eat was er test		green solution formed	
		tube. Excess aqueo ammonia was added			dark blue solution formed	

(C)	vvn	at does test (a) tell you about the type of acid in solution A?	
			[1]
(d)	(i)	Name the gas given off in test (b)(i).	[1]
	(ii)	Name the gas given off in test (b)(ii).	[,]
			[1]

	(e)	Expla	ain the observations in test (b)(iii).	
				[2]
6	The	abel	below is from a bottle of concentrated lemon drink.	
			Concentrated lemon drink	
			Ingredients: Water, sugar, citric acid, preservatives, potassium sorbate	
			(artificial sweetener). Yellow colourings E102 and E104.	
	(a)	What	t is meant by the term <i>concentrated</i> ?	
				[1]
	(b)	Predi	ict the pH of the lemon drink.	
				[1]
	(c)	Desc drink	ribe an experiment to show that two different yellow colourings are present in .	the

7 Samples of concrete were placed in solutions of different pH. The graph shows the percentage corrosion of the samples.



(a) Draw a smooth line graph on the grid [1]

(b) Which point on the grid appears to be inaccurate? Explain your reason for identifying this point.

(c) What happens to the percentage corrosion as the pH changes from 1 to 7?

[1]

8

Plan an investigation to find out which of these two oxides is the better catalyst for this decomposition. The space below can be used for a diagram.	An aqueous solution of hydrogen peroxide decomposes very slowly to form oxygen. The speed of decomposition can be increased by using a catalyst. Two possible catalysts are the solids copper(II) oxide and chromium(III) oxide.
	The space below can be used for a diagram.

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.