

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME		
CENTRE NUMBER	CANDIDATE NUMBER	
PHYSICS		0625/02

Paper 2 Core

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer all questions.

You may lose marks if you do not show your working or if you do not use appropriate units.

Take the weight of 1 kg to be 10 N (i.e. acceleration of free fall = 10 m/s^2).

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use		
1		
2		
3		
4		
5		
6		
7		
8		
9		
10		
11		
12		
Total		


May/June 2007 1 hour 15 minutes

This document consists of **19** printed pages and **1** blank page.



1 The mechanical stop-clock shown in Fig. 1.1 has

a seconds hand, which rotates once every minute and a minutes hand, which rotates once every hour.



For Examiner's Use



(a) A student uses the clock to time the intervals between trains travelling along the railway past his school.

He sets the clock to zero (both hands vertical).

As train 1 passes, he starts the clock and leaves it running.

After 35 s, train 2 passes.

On the blank face of Fig. 1.2, show the positions of the two hands of the clock as train 2 passes. Make sure it is clear which hand is which. [2]




Fig. 1.2

(b) Train 3 passes the school 4 minutes and 55 s after the clock was started.

On the blank face of Fig. 1.3, show the positions of the hands of the clock as train 3 passes. [2]



(c) Calculate the time interval between train 2 and train 3.

time interval = min s [1]

[Total: 5]






Fig. 2.1

(c) Show that the total distance travelled by the cyclist is 5400 m.

(d) Calculate the average speed of the cyclist.

average speed = m/s [2]

[Total: 11]


[4]

- sticky-tape "hinge" plank of В wood Č Fig. 3.1 (a) The cardboard is lifted as shown, using a force applied either at A or B or C. On Fig. 3.1, draw the force in the position where its value will be as small as (i) possible. [2] (ii) Explain why the position you have chosen in (a)(i) results in the smallest force. (b) Initially, the cardboard is flat on the plank of wood. A box of matches is placed on it. The cardboard is then slowly raised at the left hand edge, as shown in Fig. 3.2. stiff cardboard sticky-tape "hinge" plank of wood Fig. 3.2 State the condition for the box of matches to fall over.
- **3** A piece of stiff cardboard is stuck to a plank of wood by means of two sticky-tape "hinges". This is shown in Fig. 3.1.

For Examiner's Use

stiff cardboard

(c) The box of matches is opened, as shown in Fig. 3.3. The procedure in (b) is repeated.



4 In Fig. 4.1, a small bird, a large bird and a squirrel are on the ground under a tree.





A loud noise scares the two birds. They both fly up to the top of the tree.

- - (ii) Explain your answer to (a)(i).
 -[1]
- (b) A squirrel has the same weight as the large bird. It climbs the tree, to the same height as the birds.

How does the increase in the squirrel's gravitational potential energy compare with that of each of the two birds? Answer the question by completing the sentences below.

Compared with that of the small bird, the increase of the squirrel's potential

energy is

Compared with that of the large bird, the increase of the squirrel's potential

energy is

(c) Which creature has the least gravitational potential energy when they are at the top of the tree?

(d) The small bird flies back down to the ground.

What happens to the gravitational potential energy it had at the top of the tree?

......[2]

[2]


5 (a) Here is a list of descriptions of molecules in matter.

description		gas
free to move around from place to place		
can only vibrate about a fixed position		
closely packed		
relatively far apart		
almost no force between molecules		
strong forces are involved between molecules		

In the columns alongside the descriptions, put ticks next to those which apply to the molecules in

(i) a solid,
(ii) a gas. [4]
(b) The water in a puddle of rainwater is evaporating. Describe what happens to the molecules when the water evaporates.
[2]
[Total: 6]


6 (a) Fig. 6.1 shows how the pressure of the gas sealed in a container varies during a period of time.






Which of the following statements could explain this variation of pressure?

Tick two statements.



(b) Fig. 6.2 shows some gas trapped in a cylinder with a movable piston.






The temperature of the gas is raised.

(i) State what must happen to the piston, if anything, in order to keep the pressure of the gas constant.

[Total: 4]

7 An electric soldering iron is used to melt solder, for joining wires in an electric circuit. A soldering iron is shown in Fig. 7.1.

For Examiner's Use





Solder is a metal which melts easily. The heater coil inside the metal cylinder heats the copper tip.

(a) (i) Suggest why the tip is made of copper. (ii) Suggest why the handle is made of plastic.[1] (b) The heater coil is switched on. When the tip is put in contact with the solder, some of the heat is used to melt the solder. State the process by which the heat is transferred from the copper tip to the solder. (i) (ii) By which process or processes is the rest of the heat transferred to the surroundings? Tick the boxes alongside any of the following (you may tick as many as you think are correct). conduction convection evaporation

[2]

radiation

(c) A short time after switching on the soldering iron, it reaches a steady temperature, even though the heater coil is constantly generating heat.


The soldering iron is rated at 40W.

What is the rate at which heat is being lost from the soldering iron? Tick one box.

greater than 40W	
equal to 40W	
less than 40W	[1]

[Total: 6]

8 A square wooden block is made to rotate 3000 times per minute. A springy metal strip presses against the block, as shown in Fig. 8.1. A person nearby observes what is happening.





(a) Calculate how many times per second the block rotates.

number of rotations per second =[1]

(b) Calculate the frequency of the sound caused by this arrangement.

frequency = Hz [2]

(c) State whether or not this sound could be heard by the person nearby, and give a reason for your answer.

[1] [Total: 4]

0625/02/M/J/07

© UCLES 2007

9

(b) The circuit in Fig. 9.2 is similar to the circuit in Fig. 9.1, but it uses a resistor AB with a sliding contact.




Fig. 9.2

- (i) State the potential difference between X and Y when the sliding contact is at
 - 1. end A of the resistor, V
 - 2. end B of the resistor. V
- (ii) The sliding contact of the resistor AB is moved so that the potential difference between X and Y is 5V.

On Fig. 9.2, mark with the letter C the position of the sliding contact. [1]

[Total: 12]

[2]

For

Examiner's Use

16

11 Fig. 11.1 shows a bar magnet on a board in a region where the magnetic field of the surroundings is so weak it can be ignored. The letters N and S show the positions of the Examiner's north and south poles of the magnet. Also on the diagram are marked four dots.

Ν S

Fig. 11.1

- (a) On Fig. 11.1, carefully draw four magnetic field lines, one passing through each of the four dots. The lines you draw should begin and end either on the magnet or at the edge of the board. [5]
- (b) On one of your lines, put an arrow to show the direction of the magnetic field. [1]

[Total: 6]

For

Use

		15			
12	2 Three particles you have learned about are				
		protons, neutrons and electrons.			
	(a) I	How many of each of these particles			
		(i) are found in an α -particle,			
		number of protons =			
		number of neutrons =			
		number of electrons =[1]		
	(ii) are found in a β -particle?			
		number of protons =			
		number of neutrons =			
		number of electrons =[1]		
	(b) 🤅	Sodium-24 can be represented as $^{24}_{11}$ Na.			
How many of each of these particles are there in a neutral atom of $^{24}_{11}$ Na?					
		number of protons =			
		number of neutrons =			
		number of electrons =[3]		
		A nucleus of sodium-24 decays to become magnesium-24, by the emission of or particle. The equation below describes this change.	ne		
	-	The symbol $_{y}^{x\theta}$ represents the emitted particle.			
		$^{24}_{11}Na \rightarrow ^{24}_{12}Mg + ^{x}_{y}\theta$			
		(i) State the value of x. [1]		
	(ii) State the value of y. [1]		
	(i	ii) What type of particle is θ ?	1]		

[Total: 8]

BLANK PAGE

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.