

General Vector Space (3A)

Copyright (c) 2012 Young W. Lim.

Permission is granted to copy, distribute and/or modify this document under the terms of the GNU Free Documentation License, Version 1.2 or any later version published by the Free Software Foundation; with no Invariant Sections, no Front-Cover Texts, and no Back-Cover Texts. A copy of the license is included in the section entitled "GNU Free Documentation License".

Please send corrections (or suggestions) to youngwlim@hotmail.com.

This document was produced by using OpenOffice and Octave.

Vector Space

V : non-empty set of objects

defined operations:


addition

$$\mathbf{u} + \mathbf{v}$$

scalar multiplication

$$k \mathbf{u}$$

if the following axioms are satisfied
for all object \mathbf{u} , \mathbf{v} , \mathbf{w} and all scalar k , m



V : vector space

objects in V : vectors

1. if \mathbf{u} and \mathbf{v} are objects in V , then $\mathbf{u} + \mathbf{v}$ is in V
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
4. $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$ (zero vector)
5. $\mathbf{u} + (-\mathbf{u}) = (-\mathbf{u}) + (\mathbf{u}) = \mathbf{0}$
6. if k is any scalar and \mathbf{u} is objects in V , then $k\mathbf{u}$ is in V
7. $k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v}$
8. $(k + m)\mathbf{u} = k\mathbf{u} + m\mathbf{u}$
9. $k(m\mathbf{u}) = (km)\mathbf{u}$
10. $1(\mathbf{u}) = \mathbf{u}$

Test for a Vector Space

1. Identify the set V of objects
2. Identify the addition and scalar multiplication on V
3. Verify $u + v$ is in V and ku is in V
closure under **addition** and **scalar multiplication**
4. Confirm other axioms.

1. if u and v are objects in V , then $u + v$ is in V
2. $u + v = v + u$
3. $u + (v + w) = (u + v) + w$
4. $0 + u = u + 0 = u$ (zero vector)
5. $u + (-u) = (-u) + (u) = 0$
6. if k is any scalar and u is objects in V , then ku is in V
7. $k(u + v) = ku + kv$
8. $(k + m)u = ku + mu$
9. $k(mu) = (km)u$
10. $1(u) = u$

Subspace

a subset W of a vector space V

If the subset W is itself a vector space \Rightarrow the subset W is a **subspace** of V


1. if u and v are objects in W , then $u + v$ is in W
2. $u + v = v + u$
3. $u + (v + w) = (u + v) + w$
4. $0 + u = u + 0 = u$ (zero vector)
5. $u + (-u) = (-u) + (u) = 0$
6. if k is any scalar and u is objects in W , then ku is in W
7. $k(u + v) = ku + kv$
8. $(k + m)u = ku + mu$
9. $k(mu) = (km)u$
10. $1(u) = u$

Subspace Example (1)

In vector space R^2

any one vector	(linearly indep.)	spans R^1	line <u>through 0</u>
any two non-collinear vectors	(linearly indep.)	spans R^2	plane
any three or more vectors	(linearly dep.)	spans R^2	plane

Subspaces of R^2



Subspace Example (2)











In vector space \mathbb{R}^2

any one vector

(linearly indep.)

spans \mathbb{R}^1

line through 0



~~vector space~~

Subspace Example (3)

In vector space R^3

any one vector	(linearly indep.)	spans	R^1	line <u>through 0</u>
any two non-collinear vectors	(linearly indep.)	spans	R^2	plane <u>through 0</u>
any three vectors non-collinear, non-coplanar	(linearly indep.)	spans	R^3	3-dim space
any four or more vectors	(linearly dep.)	spans	R^3	3-dim space

Subspaces of R^3


$\{0\}$	R^1	R^2	R^3
	line <u>through 0</u>	plane <u>through 0</u>	3-dim space

Dimension


In a **finite-dimensional** vector space R^n ~~R^∞~~
all bases \rightarrow the **same number** of vectors n

many bases but the same number of basis vectors


basis $\{u_1, u_2\}$ R^2



basis $\{v_1, v_2\}$ R^2



basis $\{w_1, w_2\}$ R^2



The **dimension** of a **finite-dimensional** vector space V

$\dim(V)$



the **number** of vectors in a **basis**

Dimension of a Basis (1)

In vector space R^2

any one vector (linearly indep.) ~~spans R^2~~ line through $\mathbf{0}$

basis any two non-collinear vectors (linearly indep.) spans R^2 plane ←

any three or more vectors ~~(linearly indep.)~~ spans R^2 plane

In vector space R^3

any one vector (linearly indep.) ~~spans R^3~~ line through $\mathbf{0}$

any two non-collinear vectors (linearly indep.) ~~spans R^3~~ plane through $\mathbf{0}$

basis any three vectors non-collinear, non-coplanar (linearly indep.) spans R^3 3-dim space ←

any four or more vectors ~~(linearly indep.)~~ spans R^3 3-dim space

Dimension of a Basis (2)

In vector space R^n

any $n-1$ vectors

(linearly indep.)?

~~spans~~

~~R^n~~

line through 0

basis

n vectors of a basis

(linearly indep.)

spans

R^n

plane

any $n+1$ vectors

~~(linearly indep.)~~

spans?

R^n

plane

a finite-dimensional vector space V

a basis $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$

- a set of more than n vectors \rightarrow ~~(linearly indep.)~~
- a set of less than n vectors \rightarrow ~~spans V~~

$S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ non-empty finite set of vectors in V

S is a basis



- S linearly independent
- S spans V

Basis Test

$S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ non-empty finite set of vectors in V

S is a basis \iff $\left\{ \begin{array}{l} S \text{ linearly independent} \\ S \text{ spans } V \end{array} \right.$

V an n -dimensional vector space

$S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ a set of n vectors in V

S linearly independent \implies S is a basis

S spans V \implies S is a basis

Plus / Minus Theorem

S a nonempty set of vectors in a vector space V

S : linear independent


\mathbf{v} a vector in V but outside of $\text{span}(S)$

$\Rightarrow S \cup \{\mathbf{v}\}$: linear independent

$\mathbf{v}, \mathbf{u}_i \in S$ linear combination

$\mathbf{v} = k_1 \mathbf{u}_1 + k_2 \mathbf{u}_2 + \dots + k_n \mathbf{u}_n$

$\Rightarrow \text{span}(S) = \text{span}(S - \{\mathbf{v}\})$



Finding a Basis

S a nonempty set of vectors in a vector space V

$\left\{ \begin{array}{l} S : \text{linear independent} \\ \mathbf{v} \text{ a vector in } V \text{ but outside of } \text{span}(S) \end{array} \right. \Rightarrow S \cup \{\mathbf{v}\} : \text{linear independent}$

if S is a *linearly independent* set that is not already a basis for V ,
then S can be enlarged to a basis for V
by inserting appropriate vectors into S

$\left\{ \begin{array}{l} \mathbf{v}, \mathbf{u}_i \in S \quad \text{linear combination} \\ \mathbf{v} = k_1 \mathbf{u}_1 + k_2 \mathbf{u}_2 + \cdots + k_n \mathbf{u}_n \end{array} \right. \Rightarrow \text{span}(S) = \text{span}(S - \{\mathbf{v}\})$

if S spans V but is not a basis for V ,
then S can be reduced to a basis for V
by removing appropriate vectors from S

Vectors in a Vector Space

S a nonempty set of vectors in a vector space V

if S is a *linearly independent* set that is not already a basis for V ,
then S can be enlarged to a basis for V
by inserting appropriate vectors into S

Every *linearly independent* set in a subspace is
either a **basis** for that subspace
or can be **extended to a basis** for it

if S spans V but is not a basis for V ,
then S can be reduced to a basis for V
by removing appropriate vectors from S

Every *spanning set* for a subspace is
either a **basis** for that subspace
or has a **basis as a subset**

Dimension of a Subspace

W a subspace of a finite-dimensional vector space V

W is *finite-dimensional*

$$\dim(W) \leq \dim(V)$$

$$W = V \iff \dim(W) = \dim(V)$$

Real-Valued Functions (1)

\mathcal{V} the set of real-valued functions

defined at every x in $(-\infty, +\infty)$

$$\mathbf{u} = u(x)$$

$$\mathbf{v} = v(x)$$

$$\mathbf{u} + \mathbf{v} = u(x) + v(x)$$

$$k\mathbf{u} = ku(x)$$

1. if \mathbf{u} and \mathbf{v} are objects in \mathcal{V} , then $\mathbf{u} + \mathbf{v}$ is in \mathcal{V}
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
4. $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$ (zero vector)
5. $\mathbf{u} + (-\mathbf{u}) = (-\mathbf{u}) + (\mathbf{u}) = \mathbf{0}$
6. if k is any scalar and \mathbf{u} is objects in \mathcal{V} , then $k\mathbf{u}$ is in \mathcal{V}
7. $k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v}$
8. $(k + m)\mathbf{u} = k\mathbf{u} + m\mathbf{u}$
9. $k(m\mathbf{u}) = (km)\mathbf{u}$
10. $1(\mathbf{u}) = \mathbf{u}$

Real-Valued Functions (2)

\mathcal{V} the set of real-valued functions $\{ \sin(x), \sin(2x), \sin(3x), \dots \}$
defined at every x in $[0, 2\pi]$

$$\mathbf{u}_1 = \sin(x)$$

$$\mathbf{u}_2 = \sin(2x)$$

$$\mathbf{u}_3 = \sin(3x)$$

...

$$\mathbf{u}_m + \mathbf{v}_n = \sin(mx) + \sin(nx)$$

$$k\mathbf{u}_m = k\sin(mx)$$

1. if \mathbf{u} and \mathbf{v} are objects in \mathcal{V} , then $\mathbf{u} + \mathbf{v}$ is in \mathcal{V}
2. $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
3. $\mathbf{u} + (\mathbf{v} + \mathbf{w}) = (\mathbf{u} + \mathbf{v}) + \mathbf{w}$
4. $\mathbf{0} + \mathbf{u} = \mathbf{u} + \mathbf{0} = \mathbf{u}$ (zero vector)
5. $\mathbf{u} + (-\mathbf{u}) = (-\mathbf{u}) + (\mathbf{u}) = \mathbf{0}$
6. if k is any scalar and \mathbf{u} is objects in \mathcal{V} , then $k\mathbf{u}$ is in \mathcal{V}
7. $k(\mathbf{u} + \mathbf{v}) = k\mathbf{u} + k\mathbf{v}$
8. $(k + m)\mathbf{u} = k\mathbf{u} + m\mathbf{u}$
9. $k(m\mathbf{u}) = (km)\mathbf{u}$
10. $1(\mathbf{u}) = \mathbf{u}$

\mathcal{V} basis R^∞
linear independent

References

- [1] <http://en.wikipedia.org/>
- [2] Anton, et al., Elementary Linear Algebra, 10th ed, Wiley, 2011
- [3] Anton, et al., Contemporary Linear Algebra,