

# Contents

<b>1</b>	<b>Functions</b>	<b>2</b>
1.1	squarefree – Squarefreeness tests . . . . .	2
1.1.1	Definition . . . . .	2
1.1.2	lenstra – Lenstra’s condition . . . . .	2
1.1.3	trial_division – trial division . . . . .	3
1.1.4	trivial_test – trivial tests . . . . .	3
1.1.5	viafactor – via factorization . . . . .	3
1.1.6	viadecomposition – via partial factorization . . . . .	3
1.1.7	lenstra_ternary – Lenstra’s condition, ternary version . .	4
1.1.8	trivial_test_ternary – trivial tests, ternary version . . .	4
1.1.9	trial_division_ternary – trial division, ternary version . .	4
1.1.10	viafactor_ternary – via factorization, ternary version . . .	5

# Chapter 1

## Functions

### 1.1 squarefree – Squarefreeness tests

There are two method groups. A function in one group raises **Undetermined** when it cannot determine squarefreeness. A function in another group returns **None** in such cases. The latter group of functions have “\_ternary” suffix on their names. We refer a set  $\{\mathbf{True}, \mathbf{False}, \mathbf{None}\}$  as *ternary*.

The parameter type *integer* means either *int*, *long* or **Integer**.

This module provides an exception class.

**Undetermined** : Report undetermined state of calculation. The exception will be raised by **lenstra** or **trivial\_test**.

#### 1.1.1 Definition

We define squarefreeness as:

$n$  is squarefree  $\iff$  there is no prime  $p$  whose square divides  $n$ .

Examples:

- 0 is non-squarefree because any square of prime can divide 0.
- 1 is squarefree because there is no prime dividing 1.
- 2, 3, 5, and any other primes are squarefree.
- 4, 8, 9, 12, 16 are non-squarefree composites.
- 6, 10, 14, 15, 21 are squarefree composites.

#### 1.1.2 lenstra – Lenstra’s condition

**lenstra**( $n$ : *integer*)  $\rightarrow$  *bool*

If return value is True,  $n$  is squarefree. Otherwise, the squarefreeness is still unknown and **Undetermined** is raised. The algorithm is based on [1].

†The condition is so strong that it seems  $n$  has to be a prime or a Carmichael number to satisfy it.

Input parameter  $n$  ought to be an odd **integer**.

### 1.1.3 trial\_division – trial division

**trial\_division**( $n$ : *integer*)  $\rightarrow$  *bool*

Check whether  $n$  is squarefree or not.

The method is a kind of trial division and inefficient for large numbers.

Input parameter  $n$  ought to be an **integer**.

### 1.1.4 trivial\_test – trivial tests

**trivial\_test**( $n$ : *integer*)  $\rightarrow$  *bool*

Check whether  $n$  is squarefree or not. If the squarefreeness is still unknown, then **Undetermined** is raised.

This method do anything but factorization including Lenstra's method.

Input parameter  $n$  ought to be an odd **integer**.

### 1.1.5 viafactor – via factorization

**viafactor**( $n$ : *integer*)  $\rightarrow$  *bool*

Check whether  $n$  is squarefree or not.

It is obvious that if one knows the prime factorization of the number, he/she can tell whether the number is squarefree or not.

Input parameter  $n$  ought to be an **integer**.

### 1.1.6 viadecomposition – via partial factorization

**viadecomposition**( $n$ : *integer*)  $\rightarrow$  *bool*

Test the squarefreeness of **n**. The return value is either one of **True** or **False**; **None** never be returned.

The method uses partial factorization into squarefree parts, if such partial factorization is possible. In other cases, It completely factor **n** by trial division.

Input parameter **n** ought to be an **integer**.

### 1.1.7 `lenstra_ternary` – Lenstra’s condition, ternary version

`lenstra_ternary(n: integer) → ternary`

Test the squarefreeness of **n**. The return value is one of the ternary logical constants. If return value is **True**, **n** is squarefree. Otherwise, the squarefreeness is still unknown and **None** is returned.

†The condition is so strong that it seems **n** has to be a prime or a Carmichael number to satisfy it.

This is a ternary version of **lenstra**.

Input parameter **n** ought to be an odd **integer**.

### 1.1.8 `trivial_test_ternary` – trivial tests, ternary version

`trivial_test_ternary(n: integer) → ternary`

Test the squarefreeness of **n**. The return value is one of the ternary logical constants.

The method uses a series of trivial tests including **lenstra\_ternary**.

This is a ternary version of **trivial\_test**.

Input parameter **n** ought to be an **integer**.

### 1.1.9 `trial_division_ternary` – trial division, ternary version

`trial_division_ternary(n: integer) → ternary`

Test the squarefreeness of **n**. The return value is either one of **True** or **False**; **None** never be returned.

The method is a kind of trial division.

This is a ternary version of **trial\_division**.

Input parameter `n` ought to be an **integer**.

#### 1.1.10 `viafactor_ternary` – via factorization, ternary version

`viafactor_ternary(n: integer) → ternary`

Just for symmetry, this function is defined as an alias of **`viafactor`**.

Input parameter `n` ought to be an **integer**.

# Bibliography

- [1] H. W. Lenstra, Jr. Miller's primality test. *Information processing letters*, Vol. 8, No. 2, 1979.