

**Mathematics for natural sciences I****Exercise sheet 15****Warm-up-exercises**

EXERCISE 15.1. Show that a linear function

$$\mathbb{R} \longrightarrow \mathbb{R}, x \longmapsto ax,$$

is continuous.

EXERCISE 15.2. Prove that the function

$$\mathbb{R} \longrightarrow \mathbb{R}, x \longmapsto |x|,$$

is continuous.

EXERCISE 15.3. Prove that the function

$$\mathbb{R}_{\geq 0} \longrightarrow \mathbb{R}_{\geq 0}, x \longmapsto \sqrt{x},$$

is continuous.

EXERCISE 15.4. Let  $T \subseteq \mathbb{R}$  be a subset and let

$$f : T \longrightarrow \mathbb{R}$$

be a continuous function. Let  $x \in T$  be a point such that  $f(x) > 0$ . Prove that  $f(y) > 0$  for all  $y$  in a non-empty open interval  $]x - a, x + a[$ .

EXERCISE 15.5. Let  $a < b < c$  be real numbers and let

$$f : [a, b] \longrightarrow \mathbb{R}$$

and

$$g : [b, c] \longrightarrow \mathbb{R}$$

be continuous functions such that  $f(b) = g(b)$ . Prove that the function

$$h : [a, c] \longrightarrow \mathbb{R}$$

such that

$$h(t) = f(t) \text{ for } t \leq b \text{ and } h(t) = g(t) \text{ for } t > b$$

is also continuous.

EXERCISE 15.6. Compute the limit of the sequence

$$x_n = 5 \left( \frac{2n+1}{n} \right)^3 - 4 \left( \frac{2n+1}{n} \right)^2 + 2 \left( \frac{2n+1}{n} \right) - 3$$

for  $n \rightarrow \infty$ .

EXERCISE 15.7. Let

$$f : \mathbb{R} \longrightarrow \mathbb{R}$$

be a continuous function which takes only finitely many values. Prove that  $f$  is constant.

EXERCISE 15.8. Give an example of a continuous function

$$f : \mathbb{Q} \longrightarrow \mathbb{R},$$

which takes exactly two values.

EXERCISE 15.9. Prove that the function

$$f : \mathbb{R} \longrightarrow \mathbb{R}$$

defined by

$$f(x) = \begin{cases} x, & \text{if } x \in \mathbb{Q}, \\ 0, & \text{otherwise,} \end{cases}$$

is only at the zero point 0 continuous.

EXERCISE 15.10. Let  $T \subseteq \mathbb{R}$  be a subset and let  $a \in \mathbb{R}$  be a point. Let  $f : T \rightarrow \mathbb{R}$  be a function and  $b \in \mathbb{R}$ . Prove that the following statements are equivalent.

(1) We have

$$\lim_{x \rightarrow a} f(x) = b.$$

(2) For all  $\epsilon > 0$  there exists a  $\delta > 0$  such that for all  $x \in T$  with  $d(x, a) \leq \delta$  the inequality  $d(f(x), b) \leq \epsilon$  holds.

### Hand-in-exercises

EXERCISE 15.11. (4 points)

We consider the function

$$f(x) = \begin{cases} 1 & \text{for } x \leq -1 \\ x^2 & \text{for } -1 < x < 2 \\ -2x + 7 & \text{for } x \geq 2. \end{cases}$$

Determine the points  $x \in \mathbb{R}$  where  $f$  is continuous.

EXERCISE 15.12. (4 points)

Compute the limit of the sequence

$$b_n = 2a_n^4 - 6a_n^3 + a_n^2 - 5a_n + 3,$$

where

$$a_n = \frac{3n^3 - 5n^2 + 7}{4n^3 + 2n - 1}.$$

EXERCISE 15.13. (3 points)

Prove that the function  $f : \mathbb{R} \rightarrow \mathbb{R}$  defined by

$$f(x) = \begin{cases} 1, & \text{if } x \in \mathbb{Q}, \\ 0 & \text{otherwise,} \end{cases}$$

is for no point  $x \in \mathbb{R}$  continuous.

EXERCISE 15.14. (3 points)

Decide whether the sequence

$$a_n = \sqrt{n+1} - \sqrt{n}$$

converges and in case determine the limit.

EXERCISE 15.15. (4 points)

Determine the limit of the rational function

$$\frac{2x^3 + 3x^2 - 1}{x^3 - x^2 + x + 3}$$

at the point  $a = -1$ .